



## HOW DO CLINICIANS DIAGNOSE ATYPICAL ANOREXIA NERVOSA?

Are you a clinician currently working in the field of eating disorders, or have you done so in the past?

Atypical Anorexia Nervosa is important to understand, but it is not very clearly defined, particularly in terms of weight. Would you like to contribute to a study aiming to understand how Atypical Anorexia Nervosa can be more usefully defined?

Do you have 30 minutes to spare for this vignette study?

## WE AIM TO UNDERSTAND WHAT WEIGHT LOSS BOUNDARIES CLINICIANS USE TO DIAGNOSE SOMEONE WITH ATYPICAL ANOREXIA NERVOSA

Taking part in the study will involve reading a series of short case vignettes. After each vignette, you will be asked to rate how confident you are that the individual would be diagnosed with a number of eating disorder diagnoses (according to the DSM-5), and the treatment you would recommend. You will remain anonymous throughout.

If you would like to take part in this study, please follow the link below to the online information sheet. Once you have completed the consent form, you will be automatically directed to the survey.

[https://shef.qualtrics.com/jfe/form/SV\\_7aK\\_nz5unHGojsJ8](https://shef.qualtrics.com/jfe/form/SV_7aK_nz5unHGojsJ8)

### RESEARCHERS:

Jessica Beard, [j.beard@sheffield.ac.uk](mailto:j.beard@sheffield.ac.uk)  
Glenn Waller, [g.waller@sheffield.ac.uk](mailto:g.waller@sheffield.ac.uk)

This study has been approved by the University of Sheffield's Ethics Review Procedure.